

Durability, Degradation, Stabilization, and Lifetime Prediction (Eds.: R. L. Clough, N. C. Billingham, K. T. Gillen), ACS, Washington, DC, **1996**, pp. 59–76 (*Adv. Chem.* 249).

- [15] N. D. Searle, N. L. Maecker, L. F. E. Crewdson, *J. Polym. Sci. A* **1989**, 27, 1341.
- [16] Excitation and emission spectra of TiO₂- and CB-PBT showed only an exposure-dependent intensity increase, while TiO₂- and CB-PC and PC-PBT showed also distortions and shifts of excitation and emission spectral bands. This suggested that upon weathering the number of different types of generated fluorescent chromophores was roughly constant in PBT and was increasing in PC. For fluorescence spectra, see the Supporting Information.
- [17] Calculated for the signal-to-noise ratio of three, see: J. D. Ingle, Jr., S. R. Crouch, *Spectrochemical Analysis*, Prentice Hall, Englewood Cliffs, NJ, **1988**.
- [18] For additional experiments exploring further improvement of the screening throughput, see the Supporting Information.
- [19] Measurements of polymer fluorescence can be applicable for a variety of aromatic and other types of polymers during weathering testing. If fluorescent additives or initial impurities in polymers present a problem, molecular probes for analysis of surface oxidation products may be more applicable. See: S. R. Holmes-Farley, G. M. Whitesides, *Langmuir* **1986**, 2, 266; V. B. Ivanov, J. Behnisch, A. Hollaender, F. Mehdorn, H. Zimmermann, *Surf. Interface Anal.* **1996**, 24, 257.


A Highly Enantioselective Receptor for *N*-Protected Glutamate and Anomalous Solvent-Dependent Binding Properties**

Sara Rossi, Graham M. Kyne, David L. Turner, Neil J. Wells, and Jeremy D. Kilburn*

The development of enantioselective receptors continues to be a challenging endeavor for supramolecular chemists, and enantioselective recognition of biologically relevant molecules in competitive solvents is particularly demanding.^[1] Although numerous receptors have been developed for dicarboxylic acids and dicarboxylates,^[2] only a few enantioselective receptors for chiral dicarboxylic acids (in the neutral diprotonated form) have been described,^[3] and very few examples of enantioselective receptors for chiral dicarboxylates have been reported.^[4]

We recently described an acyclic monothiourea receptor **1a**, which bound a range of *N*-protected amino acid carboxylate salts with modest enantioselectivity.^[5] Building on this work, we have now prepared macrocyclic receptor **2**, which features two thiourea moieties flanked by carboxypyridines and separated by a chiral diamine. The receptor was designed

to produce a chiral pocket for dicarboxylates by forming up to eight hydrogen-bonding interactions with the carboxylate oxygen atoms and intramolecular hydrogen bonding with the pyridine unit to help preorganize^[6] the receptor (Scheme 1). Binding measurements, by NMR titration and isothermal




Scheme 1. Acyclic monothiourea **1** and the proposed complex of *N*-Boc-L-glutamate with macrocycle **2**. Boc = *tert*-butoxycarbonyl.

calorimetry, indicate that the macrocycle exhibits strong and highly enantioselective 1:1 binding of *N*-Boc-L-glutamate (as the dicarboxylate dianion) in relatively polar solvents (CH₃CN, DMSO) and that complexation involves a large, favorable entropic contribution. Remarkably, however, no binding of *N*-Boc-glutamate by the macrocycle is observed in a less polar solvent (CDCl₃).

The synthesis of receptor **2** was readily achieved by coupling acid **3**^[5] with (*S,S*)-1,2-diphenylethylenediamine to give bisphthalimide **4** (Scheme 2). Removal of the phthalimide protecting groups and treatment of the resulting bisamine **5** with CS₂ and DCC produced the bis(isothiocyanate) **6**. A further equivalent of bisamine **5** was added to bis(isothiocyanate) **6**, in the presence of DMAP, by syringe pump over three hours to yield the macrocyclic bistiourea **2** in 26 % yield.

The ¹H NMR spectrum^[7] of macrocycle **2** in CDCl₃ at room temperature is poorly resolved, with very broad signals for all protons, and resolution is not improved at higher temper-



Scheme 2. Reagents: a) SOCl₂; (1*S*,2*S*)-(–)-1,2-diphenylethylenediamine, DMAP, CH₂Cl₂ (68 %); b) N₂H₄·H₂O, EtOH (84 %); c) CS₂, DCC, CH₂Cl₂; d) simultaneous addition of 1 equiv of **5** and 1 equiv of **6** by syringe pumps over 3 h (26 % over steps c and d). DMAP = 4-dimethylaminopyridine, DCC = *N,N'*-dicyclohexylcarbodiimide.

[*] Prof. J. D. Kilburn, S. Rossi, G. M. Kyne, Dr. D. L. Turner, N. J. Wells
Department of Chemistry
University of Southampton
Southampton, SO17 1BJ (UK)
Fax: (+44)2380-596-805
E-mail: jdk1@soton.ac.uk

[**] We thank the Commission of the European Union (TMR Network grant "Enantioselective Separations" ERB FMRX-CT-98-0233) for financial support and postgraduate fellowships (S.R.) and the EPSRC for a quota studentship (G.M.K.).

Supporting information for this article is available on the WWW under <http://www.angewandte.org> or from the author.

atures. At -40°C , however, a well-resolved spectrum is obtained, but it is not the simple spectrum expected, given the fourfold D_2 symmetry of the macrocycle. The spectrum instead indicates twofold C_2 symmetry. In particular, the thiourea NH protons give signals at $\delta = 8.96$ and 8.08 ppm and the amide NH protons give signals at $\delta = 10.10$ and 7.88 ppm.^[7] A well-resolved and simple ^1H NMR spectrum is obtained in $[\text{D}_6]\text{DMSO}$ or CD_3CN , however, which is consistent with the expected fourfold D_2 symmetry of the macrocycle.

Binding studies^[7] with macrocycle **2** were carried out with the bis(tetrabutylammonium) salts of the enantiomers of *N*-Boc-glutamate in a range of solvents, by using standard NMR titration experiments^[8] and isothermal calorimetry.^[9,10] The addition of the *N*-Boc-L-glutamate salt to a solution of **2** in CD_3CN led to significant downfield shifts of the thiourea NH ($\Delta\delta_{\text{max}} = 1.43$ ppm), the amide NH ($\Delta\delta_{\text{max}} = 1.06$ ppm), and the benzylic CH signals ($\Delta\delta_{\text{max}} = 0.59$), which is consistent with the formation of strong hydrogen bonds and the proposed mode of binding (Scheme 1). The binding data could be readily fitted to a 1:1 binding isotherm and gave $K_a^{1:1} > 10^4 \text{ M}^{-1}$.^[11] Addition of the *N*-Boc-D-glutamate salt also led to significant downfield shifts of the NH and the benzylic CH signals. The binding data, however, could not be fitted to a 1:1 binding isotherm, but instead could be fitted to a 1:2 (host:guest) isotherm and yielded $K_a^{1:2} > 10^4 \text{ M}^{-1}$. The binding stoichiometry was confirmed by Job plots^[7,12] and is further evident from the ^1H NMR titrations, since saturation is rapidly reached after addition of approximately one equivalent of the *N*-Boc-L-glutamate salt, whereas saturation is reached only after addition of approximately two equivalents of the *N*-Boc-D-glutamate salt (Figure 1). The relatively smooth titration curve obtained with *N*-Boc-D-glutamate, and the lack of a clear transition after addition of only one equivalent of the guest, suggests that binding of the second guest molecule is stronger than the binding of the first (positive cooperativity), such that the receptor tends to saturate both binding sites simultaneously, that is, the 1:1 complex is only present in very small amounts relative to the uncomplexed receptor or the 1:2 complex.

The calorimetric binding data^[13] obtained with the *N*-Boc-L-glutamate salt as the guest confirmed the strong 1:1 binding ($K_a^{1:1} = (2.83 \pm 0.48) \times 10^4 \text{ M}^{-1}$, $\Delta G_a^{1:1} = -25.5 \pm 0.5 \text{ kJ mol}^{-1}$) which is dominated by the entropic contribution ($\Delta H_a^{1:1} = -4.5 \pm 0.2 \text{ kJ mol}^{-1}$, $T\Delta S_a^{1:1} = 21.0 \text{ kJ mol}^{-1}$). The calorimetric binding data obtained with the *N*-Boc-D-glutamate salt, using a two-site binding model, yielded a small 1:1 binding constant ($K_a^{1:1} = 38.4 \pm 1.66 \text{ M}^{-1}$, $\Delta G_a^{1:1} = -9.0 \pm 0.1 \text{ kJ mol}^{-1}$) and a large 1:2 (host:guest) binding constant ($K_a^{1:2} = (4.92 \pm 0.07) \times 10^4 \text{ M}^{-1}$, $\Delta G_a^{1:2} = -26.8 \pm 0.1 \text{ kJ mol}^{-1}$), which again is dominated by the entropic contribution ($\Delta H_a^{1:2} = -6.7 \pm 0.1 \text{ kJ mol}^{-1}$, $T\Delta S_a^{1:2} = 20.1 \text{ kJ mol}^{-1}$). This data confirms the stronger binding of the second *N*-Boc-D-glutamate guest relative to the first (positive cooperativity), as indicated by the NMR titration data, and suggests that the enantioselectivity exhibited by macrocycle **2** for the 1:1 binding of the *N*-Boc-glutamates is $> 700:1$.

Binding studies with the *N*-Boc-L-glutamate salt in $[\text{D}_6]\text{DMSO}$ give a similar picture, with significant downfield




Figure 1. Binding titration curves for macrocycle **2** showing the shift of a) the amide and b) the benzylic CH protons on addition of the bis(tetrabutylammonium) salts of *N*-Boc-L-glutamate (x) and *N*-Boc-D-glutamate (o) in CD_3CN . The graphs indicate the strong 1:1 binding with the former, and 1:2 binding with the latter (the 1:1 [host]:[guest] is reached when [guest] = 1.5 mM).

shifts of the NH signals observed on NMR titration and simple 1:1 binding, obtained by either NMR titration ($K_a^{1:1} = 3720 \text{ M}^{-1}$, $\Delta G_a^{1:1} = -20.4 \text{ kJ mol}^{-1}$) or isothermal calorimetry ($K_a^{1:1} = 2280 \pm 258 \text{ M}^{-1}$, $\Delta G_a^{1:1} = -19.2 \pm 0.3 \text{ kJ mol}^{-1}$), again with a large entropic contribution ($\Delta H_a^{1:1} = -10.7 \pm 0.3 \text{ kJ mol}^{-1}$, $T\Delta S_a^{1:1} = -8.5 \text{ kJ mol}^{-1}$). However, the binding data obtained with the *N*-Boc-D-glutamate salt in $[\text{D}_6]\text{DMSO}$ deviates from the saturation curves expected for simple 1:1 or 1:2 binding, and reliable binding constants could not be obtained. Presumably several binding equilibria (1:2, 2:1, and possibly 2:2) are competing when the *N*-Boc-D-glutamate salt is the guest, although the NMR titration binding curve approaches saturation after addition of approximately two equivalents of guest, which indicates that a 1:2 (host:guest) binding stoichiometry is dominant.^[7]

In the less competitive solvent, CDCl_3 , addition of either enantiomer of the glutamate salt did not lead to any discernible change in the ^1H NMR spectrum of the macrocycle either at -40°C or at room temperature. No change was observed even after warming the solution for several days. A 1:1 sample of macrocycle **2** and *N*-Boc-L-glutamate salt was also dissolved in CD_3CN (to allow formation of the 1:1

complex), the solvent removed, and redissolved in CDCl_3 . The resulting ^1H NMR spectrum again showed unperturbed macrocycle. Thus, it is clear that macrocycle **2** does not bind the glutamate guests in the less polar solvent (CDCl_3), and it is improbable that this is because of slow binding kinetics.

It is a generally held view that binding interactions (for example, hydrogen bonds) between polar functionalities will lead to strong complexation in a nonpolar solvent (typically CHCl_3), but to weak (or negligible) complexation in more polar (competitive) solvents or solvent mixtures, and numerous examples of this phenomenon exist. For example, *N*-tolyl-*N'*-*n*-butylurea binds tetrabutylammonium benzoate with $K_a = 1300\text{ M}^{-1}$ in CDCl_3 and $K_a = 150\text{ M}^{-1}$ in $[\text{D}_6]\text{DMSO}$,^[14] and thiourea **1a** binds the tetrabutylammonium salt of *N*-Ac-L-Phe with $K_a = 4800\text{ M}^{-1}$ in CDCl_3 and $K_a = 680\text{ M}^{-1}$ in 10% $[\text{D}_6]\text{DMSO}/\text{CDCl}_3$.^[5]

The lack of binding exhibited by macrocycle **2** in CDCl_3 , in contrast to the results observed in CH_3CN and DMSO, is thus remarkable. To probe the origin of this anomalous behavior the conformation of the macrocycle in CDCl_3 solution was established using torsion angle dynamics with NOE and scalar coupling constant constraints.^[15] The NMR studies reveal that in CDCl_3 the macrocycle adopts a tightly wrapped conformation that is stabilized by a number of intramolecular hydrogen bonds, particularly from an amide carbonyl oxygen atom to the thiourea NH and adjacent amide NH protons (Figure 2). This hydrogen-bonding motif has been previously observed in the crystal structure of **1b** which forms a dimeric pair in the solid state.^[5]




Figure 2. Conformation of macrocycle **2** in CDCl_3 as determined by ^1H NMR spectroscopy showing intramolecular H-bonding distances.

Clearly the energy required to reorganize the macrocycle into a suitable binding conformation (that is, to break the intramolecular hydrogen bonds) is not compensated for by binding interactions that would thereby be established. The more polar solvents, CH_3CN and DMSO, can solvate the hydrogen-bonding functionality of the macrocycle leading to a less rigidly constrained molecule and the ^1H NMR spectra

which reflect the fourfold D_2 symmetry of the molecule in these solvents. Solvation of the macrocycle by the more polar solvents allows tight binding of the *N*-Boc-L-glutamate salt—albeit with some conformational reorganization of the receptor (and associated energetic cost), as evidenced by the large shifts of various CH signals in the ^1H NMR spectrum on addition of the guest. Binding, in these solvents, is driven to a large extent by entropy, presumably produced by releasing bound solvent molecules from the cavity of the macrocycle. The same cavity is unable to accommodate the enantiomeric guest (*N*-Boc-D-glutamate) in a simple 1:1 binding mode, and instead binds two *N*-Boc-D-glutamate guests with a small binding constant for the first glutamate anion and a significantly larger binding constant for the second (positive cooperativity). At first sight this may seem surprising. However, the two carboxylate guests may bind on opposite faces and at opposite ends of the macrocycle, in which case the electrostatic repulsion between the two guests should be small. Furthermore, if binding of the first *N*-Boc-D-glutamate requires considerable reorganization of the receptor, and consequent energetic cost, then, once the first carboxylate is bound, the receptor may bind the second carboxylate without significant additional energetic penalty. Conformational changes in the receptor on binding *N*-Boc-D-glutamate in acetonitrile are, again, clearly evidenced by the large shifts in many of the CH signals in the ^1H NMR spectrum of the receptor (particularly the benzylic CH signals ($\Delta\delta = 0.53\text{ ppm}$)) on addition of *N*-Boc-D-glutamate.

In summary macrocycle **2** is highly enantioselective for the 1:1 binding of the *N*-protected glutamate dianion, but the binding shows anomalous solvent-dependent behavior. We are now developing analogous receptors incorporating recognition elements for the ammonium group to produce enantioselective receptors, and hence sensors, for zwitterionic glutamate.

Received: May 16, 2002 [Z19311]

- [1] For a review on enantioselective recognition, see C. S. Wilcox, *Chem. Soc. Rev.* **1993**, 383; for a recent review on synthetic receptors, see J. H. Hartley, T. D. James, C. J. Ward, *J. Chem. Soc. Perkin Trans. 1* **2000**, 3155.
- [2] For a review of carboxylate/carboxylic acid recognition, see R. J. Fitzmaurice, G. M. Kyne, D. Douheret, J. D. Kilburn, *J. Chem. Soc. Perkin Trans. 1* **2002**, 841.
- [3] a) F. Garcia-Tellado, J. Albert, A. D. Hamilton, *Chem. Commun.* **1991**, 1761; b) V. A. Montero, L. Tomlinson, K. N. Houk, F. Diederich, *Tetrahedron Lett.* **1991**, 32, 5309; c) V. Alcazar, F. Diederich, *Angew. Chem.* **1992**, 104, 1503; *Angew. Chem. Int. Ed.* **1992**, 31, 1521; d) R. J. Pieters, F. Diederich, *Chem. Commun.* **1996**, 2255; e) R. J. Pieters, J. Cuntze, M. Bonnet, F. Diederich, *J. Chem. Soc. Perkin Trans. 2* **1997**, 1891; f) J. Cuntze, F. Diederich, *Helv. Chim. Acta.* **1997**, 897; g) P. Lustenberger, E. Martinborough, T. M. Denti, F. Diederich, *J. Chem. Soc. Perkin Trans. 1* **1998**, 747; h) J. V. Hernandez, M. Almaraz, C. Raposo, M. Martin, A. Lithgow, M. Crego, C. Caballero, J. R. Moran, *Tetrahedron Lett.* **1998**, 39, 7401; i) S. Goswami, K. Ghosh, S. Dasgupta, *J. Org. Chem.* **2000**, 65, 1907.
- [4] a) J. L. Sessler, A. Andrievsky, V. Král, V. Lynch, *J. Am. Chem. Soc.* **1997**, 119, 9385; b) I. Alfonso, B. Dietrich, F. Robledo, V. Gotor, J. M. Lehn, *Helv. Chim. Acta* **2001**, 84, 280.
- [5] G. M. Kyne, M. E. Light, M. B. Hursthouse, J. de Mendoza, J. D. Kilburn, *J. Chem. Soc. Perkin Trans. 1* **2001**, 1258.
- [6] C. A. Hunter, D. H. Purvis, *Angew. Chem.* **1992**, 104, 779; *Angew. Chem. Int. Ed.* **1992**, 31, 792. See also ref [5].

- [7] Copies of the NMR spectra recorded in CDCl_3 and $[\text{D}_6]\text{DMSO}$, all binding data from NMR titration and isothermal calorimetry experiments, and Job plots are provided in the Supporting Information.
- [8] The binding constants were calculated by fitting the data using NMRtit HG software, kindly provided by Prof. C. A. Hunter, University of Sheffield. See: A. P. Bisson, C. A. Hunter, J. C. Morales, K. Young, *Chem. Eur. J.* **1998**, *4*, 845.
- [9] A. Cooper, *Curr. Opin. Chem. Biol.* **1999**, *3*, 557, and references therein.
- [10] For the specific application of isothermal calorimetry to dicarboxylate recognition, see a) B. R. Linton, M. S. Goodman, E. Fan, S. A. van Arman, A. D. Hamilton, *J. Org. Chem.* **2001**, *66*, 7313. See also, for example, b) F. P. Schmidtchen, *Org. Lett.* **2002**, *4*, 431; c) B. Linton, A. D. Hamilton, *Tetrahedron* **1999**, *55*, 6027.
- [11] The data give binding constants at the upper limit of detectability by NMR titrations (10^4 – 10^5 M^{-1}) and it can be concluded that $K_a > 10^4 \text{ M}^{-1}$, but more precise values cannot be reliably obtained from the NMR titration data: a) L. Fielding, *Tetrahedron* **2000**, *56*, 6151; b) C. S. Wilcox, *Frontiers in Supramolecular Organic Chemistry and Photochemistry* (Eds.: H.-J. Schneider, H. Dürr), VCH, Weinheim, **1991**, p. 123; c) G. Weber, S. R. Anderson, *Biochemistry* **1965**, *4*, 1942. See also the Supporting Information.
- [12] K. A. Connors, *Binding Constants*, Wiley, New York, **1987**.
- [13] The ITC data was obtained with a MicroCal VP-ITC isothermal calorimeter. Data was analyzed by nonlinear least-squares fitting using the Origin software supplied by MicroCal: T Wiseman, S. Williston, J. F. Brandts, L.-N. Lin, *Anal. Biochem.* **1989**, *179*, 131. Data was modeled using the 1:1 binding site model, or the model with two independent binding sites in cases where the data could not be fitted with the 1:1 binding site model (see ref. [10c]).
- [14] T. R. Kelly, M. H. Kim, *J. Am. Chem. Soc.* **1994**, *116*, 7072.
- [15] a) P. Güntert, C. Mumenthaler, K. Wüthrich, *J. Mol. Biol.* **1997**, *273*, 283; L. Brennan, D. L. Turner, A. C. Messias, M. L. Teodoro, J. LeGall, H. Santos, A. V. Xavier, *J. Mol. Biol.* **2000**, *298*, 61. Full details of experimental methods are provided in the Supporting Information.

Catalytic Asymmetric 1,3-Dipolar Cycloaddition Reactions of Azomethine Ylides—A Simple Approach to Optically Active Highly Functionalized Proline Derivatives**


Aase Sejr Gothelf, Kurt V. Gothelf, Rita G. Hazell, and Karl Anker Jørgensen*

The 1,3-dipolar cycloaddition reaction constitutes one of the most fundamental reactions for the stereoselective construction of five-membered heterocyclic compounds.^[1] In recent years, the development of catalytic asymmetric reactions has been one of the challenging areas within the field of 1,3-dipolar cycloaddition reactions, and especially nitrones

have been the focus of attention^[2,3] as a result of the importance of the formed isoxazolidines as building blocks for more complex molecules.

Azomethine ylides can react in a 1,3-dipolar cycloaddition reaction with alkenes to form pyrrolidines, and several examples of the formation of optically active pyrrolidines based on diastereoselective reactions are known.^[4,5] However, investigations on the metal-catalyzed enantioselective version of the 1,3-dipolar cycloaddition of azomethine ylides with alkenes are very limited. Grigg and co-workers^[6] were the first to demonstrate that applying a stoichiometric amount of chiral cobalt or manganese complexes with ephedrine derivatives as the chiral ligand could give the cycloaddition product of azomethine ylides derived from imines of glycine alkyl esters with up to 96% *ee*. It has also been mentioned that silver(I) salts in combination with chiral phosphane ligands can catalyze the 1,3-dipolar cycloaddition reaction of azomethine ylides.^[6b,7]

We present herein a new highly diastereo- and enantioselective 1,3-dipolar cycloaddition reaction of azomethine ylides with alkenes catalyzed by readily available chiral Lewis acids. Although there are few stable azomethine ylides,^[8] most are unstable. Azomethine ylides **2** can be generated from, for example, imines of glycine alkyl esters **1** by reaction with a base in the presence of a Lewis acid complex. The metal-stabilized azomethine ylide **2** reacts with an alkene **3** to give highly functionalized pyrrolidines **4** (Scheme 1).



Scheme 1. Formation of highly functionalized pyrrolidines **4** from azomethine ylides **2** and alkenes **3**. EWG = electron-withdrawing group.

The reactions of *N*-benzylidene- and *N*-(2-naphthylmethylidene) glycines **1a** and **1b**, respectively, with methyl acrylate (**3a**) and Et_3N as the base in the presence of chiral ligands such as the bisoxazolines (BOX) **5a,b**^[9] and dibenzofuranyl-2,2'-bisoxazoline (DBFOX) **5c**^[10] and Lewis acids were used in the screening process (Scheme 2). Some representative results are listed in Table 1.

The use of copper(II) salts in combination with the chiral bisoxazoline ligands **5a,b** as catalysts for the reaction of *N*-benzylidene glycinate **1a** with methyl acrylate (**3a**) gave high conversion only when using the (*S*)-*t*Bu-BOX (**5a**) ligand, but unfortunately product **4a** was formed as a racemate (Table 1, entries 1 and 2). When zinc(II)^[9e] was used as the Lewis acid instead, the reaction proceeded smoothly and **4a** was formed with 76% *ee* using **5a** as the chiral ligand and THF as the solvent at room temperature (Table 1, entry 3). Furthermore, the reaction is also highly diastereoselective as only one

[*] Prof. Dr. K. A. Jørgensen, A. S. Gothelf, Dr. K. V. Gothelf, Dr. R. G. Hazell
Danish National Research Foundation: Center for Catalysis
Department of Chemistry, Aarhus University
8000 Aarhus C (Denmark)
Fax: (+45)8619-6199
E-mail: kaj@chem.au.dk

[**] We wish to express our gratitude to The Danish National Research Foundation for financial support.

Supporting information for this article is available on the WWW under <http://www.angewandte.org> or from the author.